



Acquired Peripheral Nerve Injury Findings in Critically Ill COVID-19 Patients

CASE REPORT

LOUISE TORRES 

ÂNGELA MASSIGNAN 

RAMON GHENO

JÔNATAS FÁVERO PRIETTO DOS SANTOS 

MARCELO VIANNA RAFFO

GUILHERME JAQUET RIBEIRO 

**Author affiliations can be found in the back matter of this article*



ABSTRACT

We retrospectively analyzed clinical, NCS/EMG, and NMRI aspects of five COVID-19 intensive care unit inpatients that received mechanical ventilation. After awakening from sedation, they experienced peripheral neuromyopathic symptoms.

Teaching Point: Acquired peripheral nerve injury has been described in COVID-19 infection and knowledge of the clinical, nerve conduction studies/electromyography (NCS/EMG) and neurographic magnetic resonance imaging (NMRI) findings are crucial.

CORRESPONDING AUTHOR:

Jônatas Fávero Prietto dos Santos

Hospital Moinhos de Vento, BR
jonatas_fpbs@hotmail.com

KEYWORDS:

Covid-19; peripheral neuropathy; electromyography; MR neurography; MRI

TO CITE THIS ARTICLE:

Torres L, Massignan Â, Gheno R, dos Santos JFP, Raffo MV, Ribeiro GJ. Acquired Peripheral Nerve Injury Findings in Critically Ill COVID-19 Patients. *Journal of the Belgian Society of Radiology*. 2022; 106(1): 26, 1–4. DOI: <https://doi.org/10.5334/jbsr.2757>

